

Definition:

Foreign material or food gets stuck in the esophagus (the tube from the mouth to stomach)

Signs:

Attempting to vomit, drooling uncomfortable, and with signs often first occurring during eating

Advice:

Using an endoscope to remove the foreign body is ideal. Any perforation is life threatening

OVERVIEW

- Eating (ingestion) of foreign material or foodstuffs too large to pass through the esophagus (the tube running from the throat to the stomach), causing blockage within the open space of the tube (known as “intraluminal obstruction”)

SIGNALMENT/DESCRIPTION OF PET

Species

- Dogs
- Cats
- Dogs are more likely to have esophageal foreign bodies than are cats due to their indiscriminate eating habits

Breed Predispositions

- More common in small-breed dogs; terrier breeds tend to be more likely to have esophageal foreign bodies than other breeds

Mean Age and Range

- More common in young to middle-aged pets

SIGNS/OBSERVED CHANGES IN THE PET

- Observation of pet eating or ingesting a foreign body
- Unsuccessful attempts to vomit (known as “retching”); gagging; sluggishness (lethargy); lack of appetite (known as “anorexia”); drooling (known as “ptyalism”); regurgitation (return of food or other contents from the esophagus or stomach back up through the mouth); restlessness;

difficulty swallowing (known as “dysphagia”); and persistent gulping

- Occasionally discomfort will be noted when feeling (palpating) the neck or cranial abdomen

CAUSES

- Occurs most often with an object for which size, shape, or texture does not allow free movement through the esophagus, causing the object to become lodged before it can pass into the stomach

TREATMENT

HEALTH CARE

- Emergency care—treat as inpatients and perform an examination of the open space of the esophagus using a special instrument called an “endoscope” (general term for procedure is “endoscopy”) as soon as possible after diagnosis
- If retrieval of the foreign body using the endoscope succeeds *and* esophageal damage is minimal, the pet may be discharged the same day with no special aftercare needed

ACTIVITY

- The pet routinely may resume normal activity after a foreign body has been removed

DIET

- No change needed in most cases; may alter food to more liquid consistency in the short term
- Severe trauma to the lining of the esophagus (mucosal trauma) may require using a feeding tube to allow nutritional support during esophageal healing

SURGERY

- Endoscopy is much less traumatic and invasive than surgery
- Surgery is indicated when endoscopy fails to retrieve the foreign body; when endoscopy enables advancement of the object into the stomach, but the foreign body is too large to pass through the gastrointestinal tract; or when a large tear in the esophagus (known as “esophageal perforation”) or area of dead tissue (known as “necrosis”) requires surgical repair
- It often is less traumatic to advance a bone foreign body into the stomach than to attempt retrieval; many bone foreign bodies safely can be left to dissolve in the stomach without need for surgical removal
- Non-digestible foreign bodies advanced into the stomach may need to be removed surgically

MEDICATIONS

Medications presented in this section are intended to provide general information about possible treatment. The treatment for a particular condition may evolve as medical advances are made; therefore, the medications should not be considered as all-inclusive

In cases with significant injury to the lining of the esophagus, recommendations include the following:

- Broad-spectrum antibiotics, such as amoxicillin or Clavamox® for small perforations
- Sucralfate slurry to protect the lining of the esophagus and to allow healing
- Omeprazole or pantoprazole for suppression of stomach acid production, or for less severe cases famotidine and ranitidine may be elected
- Metoclopramide or cisapride to stimulate motility and minimize reflux esophagitis
- Local anesthetic (lidocaine gel) with water can be used to reduce esophagus pain

FOLLOW-UP CARE

PATIENT MONITORING

- Examine the esophagus closely for damage to the lining
- Mild redness (known as “erythema”) and shallow ulcers (known as “erosions”) are not uncommon following esophageal foreign body, and tend to heal uneventfully
- Survey chest x-rays (radiographs) to assess for the presence of free air in the mediastinum, the area along the midline of the chest containing the heart and other structures of the chest, other than the lungs, (free air in the mediastinum is known as “pneumomediastinum”) or in the space between the chest wall and lungs (known as “pneumothorax”)
- Monitor at least 2–3 weeks for evidence of narrowing or scarring of the esophagus (stricture formation)
- Esophageal stricture—most common clinical sign is regurgitation (return of food or other contents from the esophagus or stomach back up through the mouth); contrast x-ray (radiograph) studies of the esophagus (known as an “esophagram”) and/or follow-up endoscopic evaluation of the esophagus (known as “esophagoscopy”) may be indicated

PREVENTIONS AND AVOIDANCE

- Carefully monitor what is available in the environment (such as rocks or bones) that the pet might eat and take steps to prevent the pet from eating it
- Carefully monitor what is fed to the pet

POSSIBLE COMPLICATIONS

- Approximately 25% of pets with foreign bodies develop complications
- Complications most frequently encountered include tearing of the esophagus (esophageal perforation); narrowing or scarring of the esophagus (esophageal strictures); open tracts between the esophagus and the chest (known as “esophageal fistulas”); and severe inflammation of the esophagus (known as “esophagitis”)
- Localized, transient problems with normal movement of the esophagus can occur secondary to esophageal trauma
- Free air in the mediastinum (pneumomediastinum), free air in the space between the chest wall and lungs (pneumothorax), pneumonia, inflammation of the lining of the chest (known as “pleuritis”), inflammation of the mediastinum (known as “mediastinitis”), and tracts between the bronchus and the esophagus (known as “bronchoesophageal fistulas”) can occur secondarily to tearing or perforation of the esophagus

EXPECTED COURSE AND PROGNOSIS

- Most of these pets do well and recover uneventfully
- With complications, prognosis is guarded

KEY POINTS

- Dogs are more likely to have esophageal foreign bodies than are cats due to their indiscriminate eating habits
- Possibility of complications in approximately 25% of pets with esophageal foreign bodies
- Possibility of another esophageal foreign body (that is, the pet is a “repeat offender”)

Blackwell's Five-Minute Veterinary Consult: Canine and Feline, Sixth Edition, Larry P. Tilley and Francis W.K. Smith, Jr. © 2015 John Wiley & Sons, Inc.